

WEAK FORMULATION OF ELASTODYNAMICS IN DOMAINS WITH GROWING CRACKS

EMANUELE TASSO

ABSTRACT. In this paper we formulate and study the system of elastodynamics on domains with arbitrary growing cracks. This includes homogeneous Neumann conditions on the crack sets and mixed general Dirichlet-Neumann conditions on the boundary. The only assumptions on the crack sets are to be $(n - 1)$ -rectifiable with finite surface measure, and increasing in the sense of set inclusions. In particular they might be dense, hence the weak formulation must fall outside the usual context of Sobolev spaces and Korn's inequality. We prove existence of a solution both for the damped and undamped systems, while in the damped case we are also able to prove uniqueness and an energy balance.

Keywords: second order linear hyperbolic system, dynamic fracture mechanics, crack-ing domains, boundary conditions, bounded deformation

2010 Mathematical Subject Classification: 35L53, 35A01, 35A02, 35Q74, 26A45, 74R10.

1. INTRODUCTION

The theory of dynamic fracture mechanics contains basically three principles that can be resumed as follow

- elastodynamics off the cracks;
- energy-dissipation balance which includes also the surface energy dissipated by the crack;
- a principle dictating when a crack must grow.

For the first two conditions we refer to [15], while the third one is discussed in [16] in some more details and a maximal dissipation condition is proposed.

In this paper we focus on the first issue. Precisely we consider $\Omega \subset \mathbb{R}^n$ a regular domain as reference configuration, a *fixed* family of growing-in-time crack sets $\Gamma(t)$ contained in Ω , and $u(t, x)$ the displacement which might be essentially discontinuous for $x \in \Gamma(t)$. Then, given initial conditions, and mixed Dirichlet-Neumann boundary condition on $\partial\Omega$, we want to find a solution to the system of (possibly damped) *elastodynamics*

$$\ddot{u}(t) - \operatorname{div} [\mathbb{C} \mathcal{E}u(t)] - \gamma \operatorname{div} [\mathbb{B} \mathcal{E}\dot{u}(t)] = f(t), \text{ in } \Omega \setminus \Gamma(t) \quad (1)$$

where \mathbb{C} and \mathbb{B} are the elasticity tensor and the viscosity tensors, respectively, $\mathcal{E}u$ denotes the symmetric part of the gradient of u , div denotes the divergence operator, $f(t)$ is a vector field representing the volume force, and at each time t the system (1) is complemented with homogeneous Neumann condition on the crack

sysed

$\Gamma(t)$. This last condition reflects the fact that no external forces are acting on the crack lips. The parameter γ can take value only in $\{0, 1\}$, and in particular for $\gamma = 1$ the system is called damped, while for $\gamma = 0$ the system is called undamped.

In the corresponding quasi-static models, all the known existence results for the coupled problem $(u(t), \Gamma(t))$ without a-priori assumptions on $\Gamma(t)$, are obtained by minimizing a weak form of the Griffith's energy on function spaces with no regularity on the jump sets except the $(n-1)$ -rectifiability (see [11] [14] [10] [7]). The existence of a solution with $\Gamma(t)$ closed is obtained only in particular cases through a regularity argument (see [5] [4]). Therefore, also in the dynamic case we expect that in dealing with any general existence results, no a-priori regularity assumptions on the crack sets $\Gamma(t)$ should be assumed. For this reason we assume only that the cracks $\Gamma(t)$ are $(n-1)$ -rectifiable with finite $(n-1)$ -dimensional Hausdorff measure.

In this paper we prove that, both in the undamped and damped case, a solution actually exists.

The first issue is to give a weak formulation to the system written in (1). The presence of the cracks force at each time to solve the system on the set $\Omega \setminus \Gamma(t)$. Therefore we need to introduce suitable function spaces V_t , containing for each time t the solution $u(t)$ as well as the test functions. The scalar case, i.e. when (1) reduces to the wave equation, has been treated by Dal Maso, Larsen in [8]. Since the structure of the equation implies no bound on the amplitude of the jump of u , but only on the L^2 -norm of the gradient

$$\int_{\Omega \setminus \Gamma(t)} |\nabla u(t)|^2 dx, \quad (2)$$

appgratest

they defined the problem in the context of $GSBV(\Omega)$ (for a definition we refer to [2, Definition 4.26]). Precisely in [8] it has been shown the existence of a weak solution $u(t)$ living at each time t in the space $GSBV_2^2(\Omega; \Gamma(t))$, composed of all functions $u \in GSBV(\Omega) \cap L^2(\Omega)$ whose jump sets are contained in $\Gamma(t)$ and such that (2) is finite.

In our case the structure of the equation lead to an estimate of

$$\int_{\Omega \setminus \Gamma(t)} |\mathcal{E}u(t)|^2 dx.$$

Hence V_t needs to include all the displacements in $L^2(\Omega, \mathbb{R}^n)$ whose jump discontinuities are contained in $\Gamma(t)$ and with square integrable symmetric gradient away from the cracks. Since we assumed no regularity on the cracks, in this general context a Korn's type inequality is not true. This means that we cannot control the L^2 -norm of the gradient of $u(t)$ with the L^2 -norm of its symmetric part. As a consequence we are forced to formulate our problem in the context of BD functions, and precisely to define $V_t = GSBD_2^2(\Omega; \Gamma(t))$ (see Definition 2.1) and $V_t^* = GSBD_2^2(\Omega; \Gamma(t))^*$ its dual. Note that if $\Gamma(t)$ are closed set in Ω , then $GSBD_2^2(\Omega; \Gamma(t))$ reduces to the space of square integrable vector fields, whose symmetric gradient in the sense of distribution on $\Omega \setminus \Gamma(t)$ is square integrable.

The weak formulation of the system will be

$$\langle \ddot{u}(t), \phi \rangle_t^* + \langle \mathbb{C}\mathcal{E}u(t), \mathcal{E}\phi \rangle_{H_n} + \langle \mathbb{B}\mathcal{E}\dot{u}(t), \mathcal{E}\phi \rangle_{H_n} = \langle f(t), \phi \rangle_H \quad \forall \phi \in V_t \quad (3)$$

sysedw

for a.e. $t \in [0, T]$, where $\langle \cdot, \cdot \rangle_t^*$ denotes the duality pairing between V_t and V_t^* , $\langle \cdot, \cdot \rangle_H$ $\langle \cdot, \cdot \rangle_{H_n}$ denote the scalar product in $L^2(\Omega, \mathbb{R}^n)$ and in $L^2(\Omega, \mathbb{M}_{sym}^{n \times n})$, respectively.

We want to emphasize that the one of the most serious mathematical issues arise because these spaces are varying (increasingly) in time, so that test functions at some time t are not necessarily admissible test functions for times $s < t$. Moreover since $u(t)$ lives on each time t in different spaces V_t , we need to give a meaning to the second derivative in time $\ddot{u}(t)$ as an element of V_t^* .

While in [8] only homogeneous Neumann boundary condition was considered, in the present paper we consider also non-homogeneous mixed Dirichlet-Neumann boundary conditions on $\partial\Omega$. This introduces another difficulty when the crack sets approach the boundary, and as a consequence possible problems may occur with the boundary conditions. Indeed, when we have non-homogeneous Neumann boundary condition on a part of $\partial\Omega$, we might think that when the elastic material between this part of the boundary and the crack sets is infinitesimally small, then the elastic reaction to the traction forces will be infinitesimal too. From a mathematical point of view, the difficulties are due to the lack of continuity for the trace operator acting on functions having jump sets close to the boundary. In order to solve this problem, we make use of the results obtained in [18], which allow us to restrict our attention to a suitable space of traction forces F .

We also show an energy balance and uniqueness for the damped problem. The energy balance we are able to prove in the damped case, is a conservation of kinetic plus elastic plus dissipated energy due to the damping. For the undamped problem the energy balance, where only the kinetic plus the elastic energy are considered, is clearly false. This can be seen using the results of [3]. In the undamped case, the uniqueness is still an open problem.

This paper is organized as follows: in Section 2 we define the function spaces. In particular we define V_t , we show some functional properties about these spaces, and then we introduce the space of admissible traction forces F appearing in the Neumann part of the boundary. Then we give a precise definition of $\ddot{u}(t)$ for a.e. $t \in [0, T]$ as an element of V_t^* , and we show that under some regularity assumptions on the test functions, \ddot{u} satisfies an integration by parts formula in time.

In Section 3 we first give the definition of weak solution. Then we show an existence result for the damped equation complemented with boundary conditions, by a discrete in time approximation technique and passing to the limit when the time step goes to zero (see Theorem 3.2). More precisely to define the discrete approximate solution u_k in the time interval $(t_k^i, t_k^{i+1}]$, suppose that we have already defined u_k for $t \leq t_k^i$, and let u_k^{i+1} be the minimizer in $V_{t_{i+1}} + w(t_k^i)$ ¹ of

$$u \mapsto \left\| \frac{u - u_k^i}{\tau_k} - \frac{u_k^i - u_k^{i-1}}{\tau_k} \right\|^2 + \langle \mathbb{C} \mathcal{E}u, \mathcal{E}u \rangle + \frac{1}{\tau_k} \langle \mathbb{B} (\mathcal{E}u - \mathcal{E}u_k^i), \mathcal{E}u - \mathcal{E}u_k^i \rangle - 2 \langle f_k^i, u \rangle,$$

where $u_k^i = u_k(t_k^i)$, $\|\cdot\|$ is the norm in L^2 and $\langle \cdot, \cdot \rangle$ is the L^2 -scalar product, f_k^i is a suitable discrete approximation of f and τ_k is the time step. We define u_k on $(t_k^i, t_k^{i+1}]$ as the linear interpolation between u_k^i and u_k^{i+1} .

Then we show that the limit u of the u_k , satisfies the energy balance (38). Precisely for each k , u_k satisfies a discrete energy balance which converges to the desired energy balance for u as $k \rightarrow \infty$ (see Proposition 3.4 and Proposition 3.8). As a consequence we deduce existence and uniqueness for damped problem (see Theorem 3.7).

¹ $V_{t_{i+1}} + w(t_k^i)$ is the space of functions that jump on $\Gamma(t_{i+1})$ with Dirichlet boundary condition $w(t_k^i)$ on $\partial_D\Omega \subset \partial\Omega$.

Finally, in Section 4 we show existence of a weak solution to the undamped equation complemented with boundary conditions, always by a discrete in time approximation technique and passing to the limit when the time step goes to zero (Theorem 4.2).

Acknowledgments. The author wish to thank Prof. Gianni Dal Maso for many helpful discussions on the topic.

secno

2. NOTATION AND PRELIMINARY RESULTS

We denote the space of $n \times n$ matrices with real entries as $\mathbb{M}^{n \times n}$ endowed with the euclidean scalar product

$$\xi \cdot \eta := \sum_{j=1}^n \left(\sum_{i=1}^n \xi_{ij} \eta_{ij} \right),$$

and we denote as $|\cdot|$ the associated norm; the subspace of symmetric $n \times n$ matrices is denoted by $\mathbb{M}_{sym}^{n \times n}$. $\mathcal{L}(\mathbb{M}^{n \times n})$ is the space of continuous linear maps of $\mathbb{M}^{n \times n}$ into itself.

Let $\Omega \subset \mathbb{R}^n$ be an open set. We denote the space $L^2(\Omega, \mathbb{R}^n)$ as H , with scalar products $\langle \cdot, \cdot \rangle_H$ and with associated norm $\|\cdot\|_H$. Analogously we denote the space $L^2(\Omega, \mathbb{M}^{n \times n})$ as H_n , with scalar product $\langle \cdot, \cdot \rangle_{H_n}$ and with associated norm $\|\cdot\|_{H_n}$.

When $\mathbb{C}: \Omega \rightarrow \mathcal{L}(\mathbb{M}^{n \times n})$ is a bounded symmetric and positive definite tensor field, i.e.

- $\|\mathbb{C}\|_{L^\infty} < \infty$,
- $\mathbb{C}(x)\xi \cdot \eta = \xi \cdot \mathbb{C}(x)\eta$, $\forall \xi, \eta \in \mathbb{M}^{n \times n}$, for a.e. $x \in \Omega$,
- $\mathbb{C}(x)\xi \cdot \xi \geq \gamma_0 |\xi|^2$, $\forall \xi \in \mathbb{M}_{sym}^{n \times n}$, for a.e. $x \in \Omega$ ($\gamma_0 > 0$),

then, given any Lebesgue-measurable tensor fields $\xi: \Omega \rightarrow \mathbb{M}^{n \times n}$, in order to simplify the notation we write

$$\|\xi\|_{H_n^c} := \int_{\Omega} \mathbb{C}(x)\xi(x) \cdot \xi(x) \, dx.$$

We recall that $GSBD_2^2(\Omega)$ is the space of vector fields $u \in GSBD(\Omega)$ (see [6] for the definition of $GSBD(\Omega)$) such that $u \in L^2(\Omega, \mathbb{R}^n)$, and their symmetric approximate gradients $\mathcal{E}u$ belong to $L^2(\Omega, \mathbb{M}_{sym}^{n \times n})$.

defgsbd2

Definition 2.1. Let $\Gamma \subset \Omega$ be a countably $(\mathcal{H}^{n-1}, n-1)$ -rectifiable set (see [13, Definition 3.2.14]) with $\mathcal{H}^{n-1}(\Gamma) < \infty$. We define $GSBD_2^2(\Omega; \Gamma)$ to be the vector space of all functions $u \in GSBD_2^2(\Omega)$ whose jump sets J_u is contained in Γ . We endow $GSBD_2^2(\Omega; \Gamma)$ with the scalar product

$$\langle u, v \rangle_2 := \langle u, v \rangle_H + \langle \mathcal{E}u, \mathcal{E}v \rangle_{H_n}, \quad (4)$$

scalprod

and the induced norm will be denoted by $\|\cdot\|$.

Proposition 2.2. *The inner product space $GSBD_2^2(\Omega; \Gamma)$ is an Hilbert space.*

Proof. Suppose $(u_k)_{k \in \mathbb{N}} \subset GSBD_2^2(\Omega; \Gamma)$ is a Cauchy sequence. In particular the sequence $(u_k)_{k \in \mathbb{N}}$ converges strongly in $L^2(\Omega, \mathbb{R}^n)$ to some u . Since

$$\sup_{k \in \mathbb{N}} \left(\|u_k\|_H^2 + \|\mathcal{E}u_k\|_{H_n}^2 + \mathcal{H}^{n-1}(J_{u_k}) \right) < \infty, \quad (5)$$

by [6, Theorem 11.3], passing through a subsequence, we know that there exists $v \in GSBD(\Omega)$ such that $u_k \rightarrow v$ pointwise a.e. on Ω and $\mathcal{E}u_k \rightharpoonup \mathcal{E}v$ weakly in $L^1(\Omega, \mathbb{M}_{sym}^{n \times n})$. This implies $u = v$ and $u \in GSBD_2^2(\Omega)$. It remains to prove that

$J_u \subseteq \Gamma$.

Using [6, Theorem, 11.3], for every open set $A \subset \Omega$ we have

$$\mathcal{H}^{n-1}(J_u \cap A) \leq \liminf_{k \rightarrow \infty} \mathcal{H}^{n-1}(J_{u_k} \cap A).$$

Since the measure $\mathcal{H}^{n-1} \llcorner \Gamma$ is inner regular, for every $\epsilon > 0$ we can find a compact set $K \subset \Gamma$, such that $\mathcal{H}^{n-1}(\Gamma \setminus K) \leq \epsilon$, and so

$$\mathcal{H}^{n-1}(J_u \setminus K) \leq \liminf_{k \rightarrow \infty} \mathcal{H}^{n-1}(J_{u_k} \setminus K) \leq \liminf_{k \rightarrow \infty} \mathcal{H}^{n-1}(\Gamma \setminus K) \leq \epsilon.$$

For the arbitrariness of ϵ , we conclude that $\mathcal{H}^{n-1}(J_u \setminus \Gamma) = 0$. \square

The dual $GSBD_2^2(\Omega; \Gamma)^*$ will not be identified with the underlying Hilbert space, but instead will be endowed with a pairing consistent with the L^2 inner product, as is usually done for the duals of Sobolev spaces. Since

$$GSBD_2^2(\Omega; \Gamma) \subset L^2(\Omega, \mathbb{R}^n)$$

is a dense embedding, we have

$$L^2(\Omega, \mathbb{R}^n) = L^2(\Omega, \mathbb{R}^n)^* \subset GSBD_2^2(\Omega; \Gamma)^*,$$

and $L^2(\Omega, \mathbb{R}^n)$ is densely embedded in $GSBD_2^2(\Omega; \Gamma)^*$.

In the case Ω has also finite perimeter, the trace operator $Tr(\cdot)$ can be extended to the space $GSBD(\Omega; \Gamma)$, using the notion of *approximate limit* on the point of the reduced boundary $\mathcal{F}\Omega$ (see [18, Definition 3.9]). Moreover the following theorem holds

traceint

Theorem 2.3. *Let $\Omega \subset \mathbb{R}^n$ be an open set of finite perimeter, and let $\Gamma \subset \Omega$ be a countably $(\mathcal{H}^{n-1}, n-1)$ -rectifiable set, with $\mathcal{H}^{n-1}(\Gamma) < \infty$. Then there exists a measurable functions $\Theta: \mathcal{F}\Omega \rightarrow \mathbb{R}^+$ such that*

- (a) $\mathcal{H}^{n-1}(\{\Theta = 0\}) = 0$ and $\Theta \in L^\infty(\mathcal{F}\Omega, \mathcal{H}^{n-1})$ (in particular $\|\Theta\|_\infty \leq 1$);
- (b) For every $u \in GSBD_2^2(\Omega; \Gamma)$ we have

$$\int_{\mathcal{F}\Omega} |Tr(u)|^2 \Theta \, d\mathcal{H}^{n-1} \leq C(n, 2)(\|u\|_H + \|\mathcal{E}u\|_{H_n})^2, \quad (6)$$

tracein

where $C(n, 2)$ is a constant depending only on n and 2.

Proof. Let Θ^+ be the weight functions given by [18, Theorem 3.2], then it is enough to define

$$\Theta(x) := \Theta^+(x) \text{ for } \mathcal{H}^{n-1}\text{-a.e. } x \in \mathcal{F}\Omega,$$

then (a) and (b) are a direct consequences of [18, Theorem 3.2] with $p = 2$. \square

Now let $\partial_N \Omega$ be a Borel subset of $\mathcal{F}\Omega$. In order to impose a Neumann boundary condition in equation (1), we are led to study the continuity property of the following linear form

$$u \mapsto \int_{\partial_N \Omega} F \cdot Tr(u) \, d\mathcal{H}^{n-1} \quad u \in GSBD_2^2(\Omega; \Gamma), \quad (7)$$

linneufor

where $F: \partial_N \Omega \rightarrow \mathbb{R}^n$ is some measurable vector field. In view of inequality (6) we introduce the following spaces

Definition 2.4 (Admissible Neumann term). Let Ω , Γ , and Θ be as in the previous Theorem, and let $\partial_N \Omega$ be a Borel subset of $\mathcal{F}\Omega$. We define $N_\Theta := L^2(\partial_N \Omega, \Theta \mathcal{H}^{n-1})$,

and we denote by N_Θ^* its dual. We identify N_Θ^* with the space of measurable vector field $F: \partial_N \Omega \rightarrow \mathbb{R}^n$ such that

$$\int_{\partial_N \Omega} \frac{|F|^2}{\Theta} d\mathcal{H}^{n-1} < \infty,$$

and we consider the corresponding duality pairing between N_Θ^* and N_Θ given by

$$\langle F, g \rangle_\Theta := \int_{\partial_N \Omega} F \cdot g d\mathcal{H}^{n-1} \quad (g \in N_\Theta).$$

The induced norm is denoted by $\|\cdot\|_{N_\Theta^*}$.

Putting together the definition of N_Θ and Theorem 2.3 we have the following result

neucont

Proposition 2.5. *Let Ω , Γ , and Θ be as in Theorem 2.3. Let $\partial_N \Omega$ be a Borel subset of $\mathcal{F}\Omega$. If $F \in N_\Theta^*$ then the linear form defined in (7) belongs to $GSBD_2^2(\Omega; \Gamma)^*$.*

Proof. It is enough to use inequality (6) to have

$$\int_{\partial_N \Omega} F \cdot Tr(u) d\mathcal{H}^{n-1} \leq \|F\|_{N_\Theta^*} \|Tr(u)\|_{N_\Theta} \leq C(n, 2) \|F\|_{N_\Theta^*} (\|u\|_H + \|\mathcal{E}u\|_{H_n}).$$

□

Our choice of Neumann forces, in some sense, is natural. In fact looking at the construction of Θ made in [18], roughly speaking, it turns out that Θ measures the “closeness” of Γ to the boundary. From a physical point of view, this might be interpreted as the fact that, when the elastic material between the Neumann boundary and the crack is infinitesimally small, then the elastic reaction to the traction force will be infinitesimal too; hence, in order to reach the equilibrium, the traction forces have to decrease their intensity (proportionally to Θ).

The following Proposition is useful to prescribe a Dirichlet boundary condition on a certain portion of the boundary $\partial_D \Omega \subseteq \mathcal{F}\Omega$.

Proposition 2.6. *Let Ω , Γ be as in Proposition 2.5, let $\partial_D \Omega$ be a Borel subset of $\mathcal{F}\Omega$, and let $g: \partial_D \Omega \rightarrow \mathbb{R}$ be a measurable function. Then the set $\{u \in GSBD_2^2(\Omega; \Gamma) \mid Tr(u) = g \text{ on } \partial_D \Omega\}$ is an affine closed subspace of $GSBD_2^2(\Omega; \Gamma)$.*

Proof. The only non trivial fact is to show that it is closed with respect to the norm induced by the scalar product (4). But it is a direct consequence of [18, Theorem 4.1]. □

Definition 2.7. Let $\partial_D \Omega \subseteq \mathcal{F}\Omega$ be a Borel set. We define

$$GSBD_{2,D}^2(\Omega; \Gamma) := \{u \in GSBD_2^2(\Omega; \Gamma) \mid Tr(u) = 0, \text{ on } \partial_D \Omega\}. \quad (8)$$

Thanks to our previous proposition, $GSBD_{2,D}^2(\Omega; \Gamma)$ is actually an Hilbert space with scalar product inherited as a subspace of $GSBD_2^2(\Omega; \Gamma)$.

Now fix $T > 0$, and fix $\Gamma \subset \Omega$ a countable ($\mathcal{H}^{n-1}, n-1$)-rectifiable set with $\mathcal{H}^{n-1}(\Gamma) < \infty$. Consider for $t \in [0, T]$ an increasing family of cracks $t \mapsto \Gamma(t)$

$$\Gamma(s) \subseteq \Gamma(t) \subseteq \Gamma \text{ if } s \leq t,$$

For simplicity of notation, we denote $GSBD_2^2(\Omega; \Gamma)$ by V and $GSBD_{2,D}^2(\Omega; \Gamma(t))$ by V_t . The norm in V is denoted by $\|\cdot\|$, the norm in V_t with $\|\cdot\|_t$. Note that for $s < t$ we have $V_s \subset V_t \subset V$, and as we already mentioned, since $V \subset H$ is densely embedded in H , we have the embedding $H \subset V^*$ and the density of H in V^* . Similarly H is a

dense subspace of V_t^* for every $t \in [0, T]$. We denote the pairing between V^* and V by $\langle \cdot, \cdot \rangle$, and the associated dual norm $\|\cdot\|^*$, we denote the pairing between V_t^* and V_t by $\langle \cdot, \cdot \rangle_t$, and the associated dual norm $\|\cdot\|_t^*$. We note that these pairings are the unique continuous bilinear maps on $V^* \times V$ and $V_t^* \times V_t$ such that $\langle f, v \rangle = \langle f, v \rangle_H$ and $\langle f, v_t \rangle_t = \langle f, v_t \rangle_H$ whenever $f \in H, v \in V, v_t \in V_t$.

If $s < t$ then V_s is not dense in V_t and so V_t^* is not embedded in V_s^* . Anyway we can introduce the projection operators from V_t^* to V_s^* as the following

Definition 2.8. Let $s < t$ and let $i : V_s \rightarrow V_t$ denote the embedding $V_s \subset V_t$. Let f be an element of V_t^* . Then we define the projection map P_{st} of V_t^* onto V_s^* as

$$\langle P_{st}f, v_s \rangle_s := \langle f, i(v_s) \rangle_t \text{ for any } v_s \in V_s. \quad (9)$$

Note that the projection maps defined above are continuous and in particular $\|P_{st}f\|_s^* \leq \|f\|_t^*$. When there is no misunderstanding, we omit the notation $P_{st}f$, since the action of $f \in V_t^*$ on elements of $V_s \subset V_t$ is clear from the context.

dd **Lemma 2.9.** Let $u \in W^{1,\infty}(0, T; H)$. Assume that there exists a positive function $g \in L^2(0, T)$, such that for every $s, t \in [0, T]$ with $s < t$, we have

$$u \in W^{2,2}(t, T; V_s^*), \text{ and } \|\ddot{u}(r)\|_s^* \leq g(r) \text{ for a.e. } r \in (t, T). \quad (10) \quad \text{dd1}$$

Then there exists a set $E \subset [0, T]$ of full measure, such that for every $t \in E$ there exists $w(t) \in V_t^*$ with the following properties

$$\|w(t)\|_t^* \leq \tilde{g}(t), \quad (11) \quad \text{dd3}$$

where $\tilde{g}(t) = \limsup_{h \rightarrow 0^+} \frac{1}{h} \int_t^{t+h} g(r) dr$, and

$$\lim_{h \rightarrow 0^+} \frac{\dot{u}(t+h) - \dot{u}(t)}{h} = w(t), \text{ weakly in } V_t^*, \quad (12) \quad \text{dd4}$$

and

$$\lim_{h \rightarrow 0} \frac{\dot{u}(t+h) - \dot{u}(t)}{h} = w(t), \text{ strongly in } V_s^* \text{ for every } s < t. \quad (13) \quad \text{dd7}$$

In particular for every $s \in [0, T]$ the functions $t \mapsto u(t)$ and $t \mapsto P_{st}w(t)$, considered as a functions from (s, T) to V_s^* , belong respectively to $W^{2,2}(s, T; V_s^*)$ and $L^2(s, T; V_s^*)$, and satisfy $\ddot{u}(t) = P_{st}w(t)$ in V_s^* for a.e. $t \in (s, T)$.

wcont **Remark 2.10.** Under the previous hypothesis on u , for every $t \in [0, T]$ $\dot{u}(t)$ is a well defined element of H . More precisely the functions $\dot{u} : [0, T] \rightarrow H$ is weakly continuous, i.e. for every $t_k \rightarrow t \in [0, T]$ we have

$$\dot{u}(t_k) \rightharpoonup \dot{u}(t) \text{ weakly in } H, \text{ as } k \rightarrow \infty. \quad (14)$$

In fact thanks to our hypothesis $\dot{u}(t_k) \rightharpoonup \dot{u}(t)$ weakly in V_0^* . Moreover the sequence $(\dot{u}(t_k))_{k \in \mathbb{N}}$ is bounded in H uniformly in k , hence it is relatively sequentially compact in H . Since $H \subset V_0^*$ is an embedding, then any weak limit of $(\dot{u}(t_k))_{k \in \mathbb{N}}$ in H , must be equal to the same element $\dot{u}(t)$.

Remark 2.11. In the proof of Lemma 2.9, we are able to show that the convergence in (12) holds when we consider the incremental quotients *only for positive* h .

In the proof of Lemma 2.9 we shall use the following result on increasing sequences of subspaces of separable Hilbert spaces proved in [6, Lemma 2.3].

ihs

Lemma 2.12. *Let $\{X_t \mid t \in [0, T]\}$ be an increasing family of closed linear subspaces of a separable Hilbert space X . Then, there exists a countable set $S \subset [0, T]$ such that for all $t \in [0, T] \setminus S$, we have*

$$X_t = \overline{\bigcup_{s < t} X_s}.$$

Proof. (Lemma 2.9) Let $D \subset [0, T]$ be a countably dense set. Choose $s \in D$, then thanks to (10) for a.e. $t > s$ there exists $\dot{u}(t)$ as an element of V_s^* and $\|\dot{u}(t)\|_s^* \leq \tilde{g}(t)$. By the fact that D is countable we have a set $E' \subset (0, T)$ of full measure, such that if $t \in E'$ $\dot{u}(t)$ exists as an element of V_s^* and $\|\dot{u}(t)\|_s^* \leq \tilde{g}(t)$ for every $s \in (0, t) \cap D$. Moreover, by density, for any $s_1 < t$ there exists $s_2 \in D$ with $s_1 < s_2 < t$, and thanks to the continuity of the projection map $P_{s_1 s_2}$, we have the relation between $\dot{u}(t)$ computed in $V_{s_2}^*$ and in $V_{s_1}^*$, given by

$$\lim_{h \rightarrow 0} \frac{\dot{u}(t+h) - \dot{u}(t)}{h} = \dot{u}(t) \text{ in } V_{s_2}^* \Rightarrow \lim_{h \rightarrow 0} \frac{\dot{u}(t+h) - \dot{u}(t)}{h} = P_{s_1 s_2} \dot{u}(t) \text{ in } V_{s_1}^*, \quad (15) \quad \text{relation}$$

This means that for every $t \in E'$ and for every $s < t$

$$\dot{u}(t) \text{ exists in } V_s^*, \quad \|\dot{u}(t)\|_s^* \leq \tilde{g}(t), \quad (16) \quad \text{ubsd}$$

and precisely two derivatives computed on different V_s^* are related by (15). Hence for every $t \in E'$, $\dot{u}(t)$ is well defined as an element of V_s^* for every $s < t$.

The previous bound in (16) implies in particular

$$u \in W^{2,2}(s, T; V_s^*) \text{ for every } s \in (0, T). \quad (17)$$

Now define

$$E := E' \cup \{t \in [0, T] \mid \overline{\bigcup_{s < t} V_s} = V_t \text{ and } \tilde{g}(t) < \infty\}, \quad (18)$$

By Lemma 2.12, Lebesgue's Differentiation Theorem, and the definition of E' , it holds that E has still full measure.

Now let $t \in E$. In order to prove (11), we notice that for every $\phi \in V_t$ there exists an increasing sequence $(s_n)_{n \in \mathbb{N}}$ converging to t , and a sequence of functions $(\phi_{s_n})_{n \in \mathbb{N}}$ strongly converging to ϕ in V_t . Hence we can define $w(t): V_t \rightarrow \mathbb{R}$ as

$$\langle w(t), \phi \rangle_t := \lim_{n \rightarrow \infty} \langle \dot{u}(t), \phi_{s_n} \rangle_{s_n} \text{ for every } \phi \in V_t. \quad (19)$$

We have to show that the previous limit exists and does not depend on the approximating sequence $(\phi_{s_n})_{n \in \mathbb{N}}$. It is enough to notice that if $n > m$ then

$$\begin{aligned} \langle \dot{u}(t), \phi_{s_n} \rangle_{s_n} - \langle \dot{u}(t), \phi_{s_m} \rangle_{s_m} &= \langle \dot{u}(t), \phi_{s_n} - \phi_{s_m} \rangle_{s_n} \\ &\leq \|\dot{u}(t)\|_{s_n}^* \|\phi_{s_n} - \phi_{s_m}\|_{s_n} \\ &\leq \tilde{g}(t) \|\phi_{s_n} - \phi_{s_m}\|_t. \end{aligned} \quad (20)$$

This defines a continuous linear functional on V_t and moreover $\|w(t)\|_t^* \leq \tilde{g}(t)$. This is exactly (11).

To prove (12), we fix $\epsilon > 0$ and $\phi \in V_t$, then we can find $s < t$ and $\phi_s \in V_s$ such that $\|\phi_s - \phi\|_t \leq \epsilon$. Hence

$$\begin{aligned} \lim_{h \rightarrow 0^+} \left\langle \frac{\dot{u}(t+h) - \dot{u}(t)}{h} - w(t), \phi \right\rangle_t &= \lim_{h \rightarrow 0^+} \left\langle \frac{\dot{u}(t+h) - \dot{u}(t)}{h} - w(t), \phi_s + (\phi - \phi_s) \right\rangle_t \\ &\leq \limsup_{h \rightarrow 0^+} \frac{1}{h} \int_t^{t+h} \langle \ddot{u}(r) - w(t), \phi - \phi_s \rangle_t dr \\ &\leq \limsup_{h \rightarrow 0^+} \left(\frac{1}{h} \int_t^{t+h} g(r) dr + \tilde{g}(t) \right) \|\phi - \phi_s\|_t \\ &\leq 2\tilde{g}(t)\epsilon, \end{aligned} \tag{21}$$

where we used the fact that $u \in W^{2,2}(t, T; V_t^*)$ and the fundamental theorem of calculus. The arbitrariness of ϵ gives assertion (12) and concludes the proof. \square

lddt

Definition 2.13. Under the assumption of Lemma 2.9, the element $w(t)$ of V_t^* defined in (12) for a.e. $t \in [0, T]$ is denoted by $\ddot{u}(t)$.

intbyp

Lemma 2.14. Let u be as in Lemma 2.9 and consider $\varphi \in L^2(0, T; V) \cap W^{1,2}(0, T; H)$ such that $\varphi(t) \in V_t$ for every $t \in [0, T]$. Then the map $t \mapsto \langle \dot{u}(t), \varphi(t) \rangle_H$ is absolutely continuous on $[0, T]$ and more precisely

$$\langle \dot{u}(t_2), \varphi(t_2) \rangle_H - \langle \dot{u}(t_1), \varphi(t_1) \rangle_H = \int_{t_1}^{t_2} \langle \ddot{u}(\tau), \varphi(\tau) \rangle_\tau + \langle \dot{u}(\tau), \dot{\varphi}(\tau) \rangle_\tau d\tau, \tag{22}$$

acsprod

for every $0 \leq t_1 < t_2 \leq T$.

Proof. By Remark 2.10 we know that $t \mapsto \dot{u}(t)$ is weakly continuous in H . Therefore, since $t \mapsto \varphi(t)$ is strongly continuous in H , we deduce that $t \mapsto \langle \dot{u}(t), \varphi(t) \rangle_H$ is a continuous real valued map.

First of all we prove our assertion for $\varphi(\cdot)_h := \varphi(\cdot - h)$ instead of $\varphi(\cdot)$. Fix any $t \in [h, T-h]$. Since $\varphi(\cdot - h) \in V_t$ on the time interval $[t, t+h]$ and $u \in W^{2,2}(t, t+h; V_t^*)$, we easily deduce that

$$\langle \dot{u}(t_2), \varphi_h(t_2) \rangle_H - \langle \dot{u}(t_1), \varphi_h(t_1) \rangle_H = \int_{t_1}^{t_2} \langle \ddot{u}(\tau), \varphi_h(\tau) \rangle_\tau + \langle \dot{u}(\tau), \dot{\varphi}_h(\tau) \rangle_\tau d\tau, \tag{23}$$

acsprod1

for every $t_1, t_2 \in (t, t+h)$ ($t_1 < t_2$). Since t was arbitrary and $\langle \dot{u}(\cdot), \varphi_h(\cdot) \rangle_H$ is continuous, we can actually obtain (23) for every $t_1, t_2 \in [h, T-h]$ ($t_1 < t_2$).

Finally thanks to the fact $\varphi \in W^{1,2}(0, T; H)$ the left hand side of (23) converges to $\langle \dot{u}(t_2), \varphi(t_2) \rangle_H - \langle \dot{u}(t_1), \varphi(t_1) \rangle_H$ as $h \rightarrow 0^+$, while using also $\varphi \in L^2(0, T; V)$ (in particular the continuity of the translations in L^2), the right hand side of (23) converges to $\int_{t_1}^{t_2} \langle \ddot{u}(\tau), \varphi(\tau) \rangle_\tau + \langle \dot{u}(\tau), \dot{\varphi}(\tau) \rangle_\tau d\tau$, and we are done. \square

3. THE DAMPED SYSTEM OF ELASTODYNAMICS

From now on we consider the following standing assumptions:

- (a) $\Omega \subset \mathbb{R}^n$ is an open set of finite perimeter;
- (b) $(\Gamma(t))_{t \in [0, T]}$ is an increasing family of crack sets:

$$\Gamma(s) \subseteq \Gamma(t) \subseteq \Gamma \quad \text{for } s < t, \tag{24}$$

where $\Gamma \subseteq \Omega$ is a countably $(\mathcal{H}^{n-1}, n-1)$ -rectifiable set with $\mathcal{H}^{n-1}(\Gamma) < \infty$;

- (c) $\partial_D \Omega, \partial_N \Omega$ are two disjoint Borel subset of $\mathcal{F}\Omega$, respectively the Dirichlet and the Neumann part of the reduced boundary, such that $\partial_D \Omega \cup \partial_N \Omega = \mathcal{F}\Omega$.

(d) $\mathbb{C}, \mathbb{B} \in L^\infty(\Omega, \mathcal{L}(\mathbb{M}^{n \times n}))$ are two symmetric and uniformly elliptic tensor field i.e.:

$$\mathbb{C}(x)\xi \cdot \eta = \xi \cdot \mathbb{C}(x)\eta, \quad \mathbb{B}(x)\xi \cdot \eta = \xi \cdot \mathbb{B}(x)\eta \quad \forall \xi, \eta \in \mathbb{M}^{n \times n}, \text{ for a.e. } x \in \Omega, \quad (25)$$

and

$$\mathbb{C}(x)\xi \cdot \xi \geq \gamma_0 |\xi|^2, \quad \mathbb{B}(x)\xi \cdot \xi \geq \gamma_1 |\xi|^2, \quad \forall \xi \in \mathbb{M}_{sym}^{n \times n}, \text{ for a.e. } x \in \Omega, \quad (26) \quad \boxed{\text{ellcon}}$$

for some constants $\gamma_0, \gamma_1 > 0$.

In this section we deal with the damped system of elastodynamics:

$$\ddot{u}(t) - \operatorname{div} [\mathbb{C} \mathcal{E} u(t)] - \operatorname{div} [\mathbb{B} \mathcal{E} \dot{u}(t)] = f(t). \quad (27) \quad \boxed{\text{systdd}}$$

Now we shall give the precise definition of weak solution:

dsd

Definition 3.1. Assume (a), (b), (c) and (d). With the notation introduced in Section 2, let $f \in L^2(0, T; V^*)$, let $w \in W^{2,2}(0, T; H) \cap W^{1,2}(0, T; H^1(\Omega)^n)$ and let $F \in L^2(0, T; N_\Theta)$ where Θ is the function relative to the crack set Γ given by Theorem 2.3. We say that u is a weak solution to (27) on the time dependent domain $t \mapsto \Omega \setminus \Gamma(t)$ with Dirichlet boundary condition $w(t)$ on $\partial_D \Omega$, Neumann boundary condition $F(t)$ on $\partial_N \Omega$, and homogeneous Neumann boundary conditions on $\Gamma(t)$, if

$$u \in W^{1,\infty}(0, T; H) \cap W^{1,2}(0, T; V). \quad (28) \quad \boxed{\text{ds2dd}}$$

$$\text{For every } t \in [0, T] \quad u(t) - w(t) \in V_t. \quad (29) \quad \boxed{\text{ds3dd}}$$

$$\text{For every } s \in [0, T] \quad u \in W^{2,2}(s, T; V_s^*), \text{ and} \quad (30) \quad \boxed{\text{ds4dd}}$$

$$\|P_{st} \ddot{u}(t)\|_s^* \leq g(t) \text{ for a.e. } t \in (s, T), \text{ for some } g \in L^2(0, T). \quad (31) \quad \boxed{\text{ds5dd}}$$

$$\lim_{h \rightarrow 0^+} \int_h^T \frac{\|\dot{u}(t) - \dot{u}(t-h)\|_H^2}{h} dt = 0. \quad (32) \quad \boxed{\text{ds6dd}}$$

For a.e. $t \in [0, T]$

$$\begin{aligned} & \langle \ddot{u}(t), \phi \rangle_t + \langle \mathbb{C} \mathcal{E} u(t), \mathcal{E} \phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E} \dot{u}(t), \mathcal{E} \phi \rangle_{H_n} - \langle F(t), \operatorname{Tr}(\phi) \rangle_\Theta = \\ & = \langle f(t), \phi \rangle_t, \text{ for every } \phi \in V_t \end{aligned} \quad (33) \quad \boxed{\text{ds7dd}}$$

where $\ddot{u}(t)$ is the one given by Definition 2.13.

Given $u^0 \in V_0$ and $u^1 \in H$, since $t \mapsto u(t)$ is strongly continuous in V the initial value for u^0 is well defined as element of V_0 . Moreover we are able to prescribe the initial conditions for $\dot{u}(0)$ asking

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_0^h \|\dot{u}(t) - u^1\|_H^2 dt = 0. \quad (34) \quad \boxed{\text{incond}}$$

The work of the external forces on the solution u over a time interval $[t_1, t_2] \subset [0, T]$ is given by

$$\mathcal{W}_{load}(u; t_1, t_2) := \int_{t_1}^{t_2} \langle f(t), \dot{u}(t) \rangle_t dt \quad (35)$$

which is well defined by (28) and the fact $f \in L^2(0, T; V^*)$. One would expect that the work on the solution u due to the varying Dirichlet boundary conditions w and Neumann boundary conditions F over a time interval $[t_1, t_2] \subset [0, T]$ is given by

$$\begin{aligned} & \mathcal{W}_{bdry}(u; t_1, t_2) := \mathcal{W}_{bdry}^D(u; t_1, t_2) + \mathcal{W}_{bdry}^N(u; t_1, t_2) \\ & := \int_{t_1}^{t_2} \left(\int_{\partial_D \Omega} \mathcal{E} u(t) \nu \cdot \dot{w}(t) d\mathcal{H}^{n-1} \right) dt + \int_{t_1}^{t_2} \left(\int_{\partial_N \Omega} F(t) \cdot \dot{u}(t) d\mathcal{H}^{n-1} \right) dt. \end{aligned} \quad (36)$$

Unfortunately, under the assumptions (28)-(31) the trace of the normal derivative $\partial_\nu u(t)$ cannot be defined, not even in a weaker sense, because $\mathcal{E}u(t)$ in general belongs only to H_n . We decide to solve this problem following [9, Proposition 3.1], by using the weak formulation of the work due to the Dirichlet boundary conditions:

$$\begin{aligned} & \mathcal{W}_{bdry}^D(u; t_1, t_2) := \langle \dot{u}(t_2), \dot{u}(t_1) \rangle_H - \langle \dot{u}(t_1), \dot{w}(t_1) \rangle_H \\ & - \int_{t_1}^{t_2} \langle \ddot{w}(t), \dot{u}(t) \rangle_H dt - \int_{t_1}^{t_2} \langle f(t), \dot{w}(t) \rangle_t dt + \int_{t_1}^{t_2} \langle \mathbb{C} \mathcal{E}u(t) + \mathbb{B} \mathcal{E}\dot{u}(t), \mathcal{E}\dot{w}(t) \rangle_{H_n} dt. \end{aligned} \quad (37)$$

With these notations, the energy balance that we are able to prove for the solution u to (27) has the following form:

$$\begin{aligned} & \frac{1}{2} \|\dot{u}(t)\|_H^2 + \frac{1}{2} \|\mathcal{E}u(t)\|_{H_n^c}^2 + \frac{1}{2} \int_0^t \frac{1}{2} \|\mathcal{E}\dot{u}(\tau)\|_{H_n^{\mathbb{B}}}^2 d\tau = \\ & = \frac{1}{2} \|u^1\|_H^2 + \frac{1}{2} \|\mathcal{E}u^0\|_{H_n^c}^2 + \mathcal{W}_{load}(u; 0, t) + \mathcal{W}_{bdry}(u; 0, t). \end{aligned} \quad (38) \quad \boxed{\text{enbaldw}}$$

Now we prove the main result:

mresd

Theorem 3.2 (Partial existence). *Assume (a), (b), (c) and (d). Let f , w and F be as in Definition 3.1. Then, given two initial conditions $u^0 \in V_0$ and $u^1 \in H$, there exists a function u satisfying (28)-(31) and (33) of Definition 4.1, with initial conditions $u(0) = u^0$ and (34). Moreover u satisfies the energy inequality*

$$\begin{aligned} & \frac{1}{2} \|\dot{u}(t)\|_H^2 + \frac{1}{2} \|\mathcal{E}u(t)\|_{H_n^c}^2 + \frac{1}{2} \int_0^t \frac{1}{2} \|\mathcal{E}\dot{u}(\tau)\|_{H_n^{\mathbb{B}}}^2 d\tau \leq \\ & \leq \frac{1}{2} \|u^1\|_H^2 + \frac{1}{2} \|\mathcal{E}u^0\|_{H_n^c}^2 + \mathcal{W}_{load}(u; 0, t) + \mathcal{W}_{bdry}(u; 0, t) \end{aligned} \quad (39) \quad \boxed{\text{enined}}$$

for a.e. $t \in [0, T]$.

Remark 3.3. Since $F \in L^2(0, T; N_\Theta)$, by Proposition 2.5 we have that $\langle F(t), Tr(\phi) \rangle_\Theta$ is actually a duality pairing between V_t^* and V_t . Therefore we can absorb the Neumann term into the forcing term defining

$$\langle \tilde{f}(t), \phi \rangle_t := \langle f(t), \phi \rangle_t + \langle F(t), Tr(\phi) \rangle_\Theta,$$

and we can reduce ourselves to prove the Theorem when (33) has the simplest form

$$\langle \ddot{u}(t), \phi \rangle_t + \langle \mathbb{C} \mathcal{E}u(t), \mathcal{E}\phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E}\dot{u}(t), \mathcal{E}\phi \rangle_{H_n} = \langle f(t), \phi \rangle_t.$$

Proof. For $k \in \mathbb{N}$, we set $\tau_k := T/k$ and $t_k^j := j\tau_k$. For $j = 1, 2, \dots, k$ we define $f_k^j \in V_T^*$ by

$$f_k^j := \frac{1}{\tau_k} \int_{t_k^j}^{t_k^{j+1}} f(\tau) d\tau, \quad (40)$$

and

$$w_k^j := w(t_k^j), \quad (41)$$

(we use $w \in W^{1,2}(0, T; H^1(\Omega)^n)$ implies that for every $t \in [0, T]$ $w(t)$ is well defined in $H^1(\Omega)^n$).

Inductively we define u_k^j for $j = -1, 0, \dots, k$ by the following:

$$u_k^0 := u^0, \quad u_k^{-1} := u^0 - \tau_k u^1; \quad (42) \quad \boxed{\text{condind}}$$

then, for $j = 0, 1, \dots, k-1$, the function u_k^{j+1} is the minimizer in $V_{t_k^{j+1}} + w_k^j$ of

$$u \mapsto \left\| \frac{u - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k} \right\|_H^2 + \|\mathcal{E}u\|_{H_n^c}^2 + \frac{1}{\tau_k} \left(\|\mathcal{E}u - \mathcal{E}u_k^j\|_{H_n}^2 - 2\langle f_k^j, u \rangle_{t_k^{j+1}} \right). \quad (43)$$

Thanks to the ellipticity hypothesis on \mathbb{C} and \mathbb{B} , at each step the above functional is coercive in $V_{t_k^{j+1}} + w_k^j$ because it is greater than

$$c_k \left[\|u\|_H^2 + (\gamma_0 + \gamma_1) \|\mathcal{E}u\|_{H_n}^2 \right] - 2\|f_k^j\|_{t_k^{j+1}}^* (\|u\|_H + \|\mathcal{E}u\|_{H_n}) - a_k^{j+1} \quad (44)$$

where $c_k := \min\{1, 1/\tau_k^2\}$, a_k^{j+1} is a constant depending only on k, j , and $\gamma_0, \gamma_1 > 0$ are the ellipticity constants introduced in (26). Writing the Euler equation for the minimizer u_k^{j+1} we have

$$\left\langle \frac{u_k^{j+1} - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k}, \frac{\phi}{\tau_k} \right\rangle_H + \langle \mathbb{C} \mathcal{E}u_k^{j+1}, \mathcal{E}\phi \rangle_{H_n} + \frac{1}{\tau_k} \langle \mathbb{B}(\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j), \mathcal{E}\phi \rangle_{H_n} = \langle f_k^j, \phi \rangle_{t_k^{j+1}} \quad (45)$$

disped

for every $\phi \in V_{t_k^{j+1}}$. Then using $u_k^{j+1} - u_k^j - (w_k^{j+1} - w_k^j) \in V_{t_k^{j+1}}$ as ϕ we can write

$$\begin{aligned} & \left\| \frac{u_k^{j+1} - u_k^j}{\tau_k} \right\|_H^2 - \left\langle \frac{u_k^{j+1} - u_k^j}{\tau_k}, \frac{u_k^j - u_k^{j-1}}{\tau_k} \right\rangle_H - \left\langle \frac{u_k^{j+1} - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k}, \frac{w_k^{j+1} - w_k^j}{\tau_k} \right\rangle_H \\ & + \|\mathcal{E}u_k^{j+1}\|_{H_n^c}^2 - \langle \mathbb{C} \mathcal{E}u_k^{j+1}, \mathcal{E}u_k^j \rangle_{H_n} - \langle \mathbb{C} \mathcal{E}u_k^{j+1}, \mathcal{E}w_k^{j+1} - \mathcal{E}w_k^j \rangle_{H_n} + \frac{1}{\tau_k} \|\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j\|_{H_n}^2 \\ & - \frac{1}{\tau_k} \langle \mathbb{B}(\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j), \mathcal{E}w_k^{j+1} - \mathcal{E}w_k^j \rangle_{H_n} = \langle f_k^j, u_k^{j+1} - u_k^j \rangle_{t_k^{j+1}} - \langle f_k^j, w_k^{j+1} - w_k^j \rangle_{t_k^{j+1}}. \end{aligned} \quad (46)$$

Now using the identity $\|a\|^2 - \langle a, b \rangle = \frac{1}{2}\|a\|^2 + \frac{1}{2}\|a - b\|^2 - \frac{1}{2}\|b\|^2$, multiplying by 2, and rearranging, we get

$$\begin{aligned} & \left\| \frac{u_k^{j+1} - u_k^j}{\tau_k} \right\|_H^2 + \left\| \frac{u_k^{j+1} - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k} \right\|_H^2 - 2 \left\langle \frac{u_k^{j+1} - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k}, \frac{w_k^{j+1} - w_k^j}{\tau_k} \right\rangle_H \\ & + \|\mathcal{E}u_k^{j+1}\|_{H_n^c}^2 + \|\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j\|_{H_n^c}^2 + 2\langle \mathbb{C} \mathcal{E}u_k^{j+1}, \mathcal{E}w_k^{j+1} - \mathcal{E}w_k^j \rangle_{H_n} + \\ & + 2\frac{1}{\tau_k} \|\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j\|_{H_n}^2 - 2\frac{1}{\tau_k} \langle \mathbb{B}(\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j), \mathcal{E}w_k^{j+1} - \mathcal{E}w_k^j \rangle_{H_n} = \\ & = \left\| \frac{u_k^j - u_k^{j-1}}{\tau_k} \right\|_H^2 + \|\mathcal{E}u_k^j\|_{H_n^c}^2 + 2\langle f_k^j, u_k^{j+1} - u_k^j \rangle_{t_k^{j+1}} - 2\langle f_k^j, w_k^{j+1} - w_k^j \rangle_{t_k^{j+1}}. \end{aligned} \quad (47)$$

Summing from $j = 0$ to $i \in \{1, \dots, k\}$ and using (42), we get

$$\begin{aligned}
& \left\| \frac{u_k^{i+1} - u_k^i}{\tau_k} \right\|_H^2 + \|\mathcal{E}u_k^{i+1}\|_{H_n^c}^2 + \sum_{j=0}^i \left\| \frac{u_k^{j+1} - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k} \right\|_H^2 + \sum_{j=0}^i \|\mathcal{E}u_n^{j+1} - \mathcal{E}u_k^j\|_{H_n^c}^2 + \\
& + 2 \frac{1}{\tau_k} \sum_{j=0}^i \|\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j\|_{H_n}^2 = \|u^1\|_H^2 + \|\mathcal{E}u^0\|_{H_n^c}^2 + 2 \sum_{j=0}^i \langle \langle f_k^j, u_k^{j+1} - u_k^j \rangle \rangle_{t_k^{j+1}} + \\
& + 2 \sum_{j=0}^i \langle \langle f_k^j, w_k^{j+1} - w_k^j \rangle \rangle_{t_k^{j+1}} + 2 \sum_{j=0}^i \langle \langle \mathbb{C} \mathcal{E}u_k^{j+1} + \frac{1}{\tau_k} \mathbb{B}(\mathcal{E}u_k^{j+1} - \mathcal{E}u_k^j), \mathcal{E}w_k^{j+1} - \mathcal{E}w_k^j \rangle \rangle_{H_n} + \\
& - 2 \sum_{j=0}^i \left\langle \frac{u_k^j - u_k^{j-1}}{\tau_k}, \frac{w_k^{j+1} - w_k^j}{\tau_k} - \frac{w_k^j - w_k^{j-1}}{\tau_k} \right\rangle_H + 2 \left\langle \frac{u_k^{i+1} - u_k^i}{\tau_k}, \frac{w_k^{i+1} - w_k^i}{\tau_k} \right\rangle_H - 2 \left\langle \frac{u_k^1 - u_k^0}{\tau_k}, \frac{w_k^1 - w_k^0}{\tau_k} \right\rangle_H
\end{aligned} \tag{48}$$

discen1d

We define the piece-wise affine discrete approximations $u_k, v_k, w_k, z_k: [0, T] \rightarrow V$ for $t \in (t_k^j, t_k^{j+1}]$ by

$$u_k(t) := u_k^j + \frac{t - t_k^j}{\tau_k} (u_k^{j+1} - u_k^j), \tag{49} \quad \text{discs1d}$$

$$w_k(t) := w_k^j + \frac{t - t_k^j}{\tau_k} (w_k^{j+1} - w_k^j), \tag{50} \quad \text{discs4d}$$

$$v_k(t) := \frac{u_k^j - u_k^{j-1}}{\tau_k} + \frac{t - t_k^j}{\tau_k} \left(\frac{u_k^{j+1} - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k} \right) \tag{51} \quad \text{discs3d}$$

$$z_n(t) := \frac{w_k^j - w_k^{j-1}}{\tau_k} + \frac{t - t_k^j}{\tau_k} \left(\frac{w_k^{j+1} - w_k^j}{\tau_k} - \frac{w_k^j - w_k^{j-1}}{\tau_k} \right), \tag{52} \quad \text{discs5d}$$

and the piece-wise constant discrete approximations $\tilde{u}_k, \tilde{w}_k, f_k: [0, T] \rightarrow V$ for $t \in (t_k^j, t_k^{j+1}]$ by

$$\tilde{u}_k(t) := u_k^{j+1}, \tilde{w}_k(t) := w_k^{j+1}, f_k(t) := f_k^j. \tag{53} \quad \text{discs2d}$$

(54)

Rewriting (48) in terms of $u_k, w_k, \tilde{u}_k, v_k, z_k$ we get the discrete energy balance for every $t \in (t_k^j, t_k^{j+1})$:

$$\begin{aligned}
& \|\dot{u}_k(t)\|_H^2 + \|\mathcal{E}u_k(t_k^{j+1})\|_{H_n^c}^2 + \tau_k \int_0^{t_k^{j+1}} \|\dot{v}_k(\tau)\|_H^2 d\tau + \tau_k \int_0^{t_k^{j+1}} \|\mathcal{E}\dot{u}_k(\tau)\|_{H_n^c}^2 d\tau + \\
& + \int_0^{t_k^{j+1}} \|\mathcal{E}\dot{u}_k(\tau)\|_{H_n^b}^2 d\tau = \|u^1\|_H^2 + \|\mathcal{E}u^0\|_{H_n^c}^2 + 2 \int_0^{t_k^{j+1}} \langle \langle f_k(\tau), \dot{u}_k(\tau) \rangle \rangle_{t_k^{j+1}} d\tau + \\
& + 2 \int_0^{t_k^{j+1}} \langle \langle f_k(\tau), \dot{w}_k(\tau) \rangle \rangle_{t_k^{j+1}} + \langle \langle \mathbb{C} \mathcal{E}\tilde{u}_k(\tau), \mathcal{E}\dot{w}_k(\tau) \rangle \rangle_{H_n} + \langle \langle \mathbb{B} \mathcal{E}\dot{u}_k(\tau), \mathcal{E}\dot{w}_k(\tau) \rangle \rangle_{H_n} d\tau + \\
& - 2 \int_0^{t_k^{j+1}} \langle \langle \dot{u}_k(\tau), \dot{z}_k(\tau) \rangle \rangle_H d\tau + 2 \langle \langle \dot{u}_k(t), \dot{w}_k(t) \rangle \rangle_H - 2 \langle \langle u^1, \dot{w}_k(0) \rangle \rangle_H
\end{aligned} \tag{55}$$

discen2d

Let $M_k := \sup_{t \in (0, T)} \|\dot{u}_k(t)\|_H$, $L_k := \sup_{t \in (0, T)} \|\mathcal{E}\tilde{u}_k(t)\|_{H_n^c}$. By (55) we can give the estimate

$$M_k^2 + L_k^2 + \|\mathcal{E}\dot{u}_k\|_{L^2(0, T; H_n^b)}^2 \leq a(M_k + L_k + \|\mathcal{E}\dot{u}_k\|_{L^2(0, T; H_n^b)}) + b, \tag{56}$$

where a and b are constants that depends only on $\|f\|_{L^2(0,T;V^*)}$, $\|w\|_{W^{1,2}(0,T;V)}$, $\|w\|_{W^{2,2}(0,T;H)}$, $\|u_1\|_H$ and on T . As a consequence we can deduce the followings

$$\mathcal{E}u_k(t) \text{ and } \mathcal{E}\tilde{u}_k(t) \text{ are bounded in } H_n \text{ uniformly in } t \text{ and } k, \quad (57) \quad \boxed{\text{bpds1d}}$$

$$\dot{u}_k(t) \text{ and } v_k(t) \text{ are bounded in } H \text{ uniformly in } t \text{ and } k \quad (58) \quad \boxed{\text{bpds3d}}$$

$$\mathcal{E}\dot{u}_k \text{ is bounded in } L^2(0,T;H_n) \text{ uniformly in } k. \quad (59) \quad \boxed{\text{bpds2d}}$$

Note also that with the fact $u^0 \in H$ implies that u_k is bounded in H uniformly in t and k . This with (57) gives

$$u_k(t) \text{ is bounded in } V \text{ uniformly in } t \text{ and } k. \quad (60) \quad \boxed{\text{bpds4d}}$$

Furthermore, using (49)-(51) and (53), we can rewrite (45) for all $t \in (t_k^j, t_k^{j+1})$ as

$$\langle \dot{v}_k(t), \phi \rangle_H + \langle \mathbb{C} \mathcal{E}\tilde{u}_k(t), \mathcal{E}\phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E}\dot{u}_k(t), \mathcal{E}\phi \rangle_{H_n} = \langle f_k(t), \phi \rangle_{t_k^{j+1}} \quad (61) \quad \boxed{\text{dispde1d}}$$

for every $\phi \in V_{t_k^{j+1}}$. The last equation leads us to write for all $t \in (t_k^j, t_k^{j+1})$

$$\|\dot{v}_k(t)\|_{t_k^{j+1}}^* \leq \|\mathcal{E}\tilde{u}_k(t)\|_{H_n^c} + \|\mathcal{E}\dot{u}_k(t)\|_{H_n} + \|f_k(t)\|_{t_k^{j+1}}^*. \quad (62) \quad \boxed{\text{bpds5d}}$$

In particular fix $s \in [0, T]$, then for every $t_1, t_2 \in [s, T]$ with $t_1 < t_2$, we have

$$\int_{t_1}^{t_2} \|P_{st}\dot{v}_k(t)\|_s^* dt \leq \int_{t_1}^{t_2} (\|\mathcal{E}\tilde{u}_k(t)\|_{H_n^c} + \|\mathcal{E}\dot{u}_k(t)\|_{H_n} + \|f_k(t)\|_s^*) dt. \quad (63) \quad \boxed{\text{acestd}}$$

Using (57)-(59), and (62), there exist a constant M such that eventually passing through a subsequence, if we call v a weak limit of v_k in $W^{1,2}(s, T; V_s^*)$, and \tilde{g} a weak limit of $t \mapsto \|\mathcal{E}\dot{u}_k(t)\|_{H_n}$ in $L^2(0, T)$, we have

$$\int_{t_1}^{t_2} \|P_{st}\dot{v}(t)\|_s^* d\tau \leq M|t_2 - t_1| + \int_{t_1}^{t_2} (\tilde{g}(t) + \|f(t)\|_s^*) dt, \quad (64)$$

for every $t_1, t_2 \in [s, T]$ with $t_1 < t_2$.

Now if we fix a dense set $D \subset [0, T]$, using a diagonal argument we obtain a subsequence, not relabeled, such that

$$u_k \rightharpoonup u \text{ weakly in } W^{1,2}(0, T; V) \quad (65) \quad \boxed{\text{estlem3d}}$$

$$v_k \rightharpoonup v \text{ weakly in } L^2(0, T; H), \quad (66) \quad \boxed{\text{estlem2d}}$$

$$v_k \rightharpoonup v \text{ weakly in } W^{1,2}(s, T; V_s^*) \text{ for every } s \in D, \quad (67) \quad \boxed{\text{estlem1d}}$$

and

$$\int_{t_1}^{t_2} \|P_{st}\dot{v}(t)\|_s^* d\tau \leq M|t_2 - t_1| + \int_{t_1}^{t_2} (\tilde{g}(t) + \|f(t)\|_s^*) dt, \quad (68) \quad \boxed{\text{estlem6d}}$$

for every $t_1, t_2 \in [s, T]$ with $t_1 < t_2$.

Moreover using the continuity of the projection maps P_{st} , it follows that (67) and (68) become

$$v_k \rightharpoonup v \text{ weakly in } W^{1,2}(s, T; V_s^*) \text{ for every } s \in [0, T], \quad (69) \quad \boxed{\text{estlem5d}}$$

and

$$\int_{t_1}^{t_2} \|P_{st}\dot{v}(t)\|_s^* d\tau \leq M|t_2 - t_1| + \int_{t_1}^{t_2} (\tilde{g}(t) + \|f(t)\|_s^*) dt, \quad (70) \quad \boxed{\text{estlem4d}}$$

for every $s \in [0, T]$ and every $t_1, t_2 \in [s, T]$ with $t_1 < t_2$. In particular

$$\|P_{st}\dot{v}(t)\|_s^* \leq M + \tilde{g}(t) + \|f(t)\|_s^*, \quad (71) \quad \boxed{\text{pwestd}}$$

for every $s \in [0, T]$ and a.e. $t > s$.

By (58) it is easy to see that in fact

$$u \in W^{1,\infty}(0, T; H), \quad (72)$$

and this with convergence (65) gives (28).

Now we want to show

$$\dot{u}(t) = v(t) \text{ in } H \text{ for a.e. } t \in [0, T]. \quad (73)$$

First of all for $t \in (t_k^j, t_k^{j+1})$ we have $\dot{u}_k(t) = v_k(t_k^{j+1})$ and so

$$\|\dot{u}_k(t) - v_k(t)\|_{t_k^{j+1}}^* = \|v_k(t_k^{j+1}) - v_k(t)\|_{t_k^{j+1}}^* \leq \int_{t_k^j}^{t_k^{j+1}} \|\dot{v}_k(\tau)\|_{t_k^{j+1}}^* d\tau \leq \tau_k^{\frac{1}{2}} C, \quad (74)$$

where C is a uniform bound on the L^2 norm of the right-hand side of (62). Then for all $s < t$ we have $\|\dot{u}_k(t) - v_k(t)\|_s^* \leq \tau_k^{\frac{1}{2}} C$, and this together with (66) implies $\dot{u}_k \rightharpoonup v$ weakly in $L^2(s, T; V_s^*)$ for any $s \in [0, T)$. But also $\dot{u}_k \rightharpoonup \dot{u}$ weakly in $L^2(0, T; H)$ by (65). So $v(t) = \dot{u}(t)$ in V_s^* , for every $s \in [0, T)$ and for a.e. $t \in (s, T)$. Since $v(t)$ and $\dot{u}(t)$ belong to H , and H is embedded in V_s^* for every $s \in [0, T]$ we finally get that $v(t) = \dot{u}(t)$ as elements of H for a.e. $t \in [0, T]$. This together with (65) and (70) allows to conclude that

$$u \in W^{2,2}(t, T; V_s^*) \text{ for every } s, t \in [0, T] \text{ with } s < t. \quad (75) \quad \boxed{\text{alem1d}}$$

Let $g(t) := M + \tilde{g}(t) + \|f(t)\|_t^*$ then by (71) we have

$$\|P_{st}\ddot{u}(t)\|_s^* \leq g(t), \quad (76) \quad \boxed{\text{alem2d}}$$

for every $s \in [0, T)$ and for a.e. $t > s$.

Now we investigate the convergence of the constant piecewise interpolated \tilde{u}_k . Since by (65) u_k are Lipschitz with values in H uniformly in k , as before we get that

$$\tilde{u}_k \rightharpoonup u \text{ weakly in } L^2(0, T; H), \quad (77)$$

and since by (57) $\mathcal{E}\tilde{u}_k$ is bounded in $L^2(0, T; H_n)$, we also obtain that up to subsequences

$$\tilde{u}_k \rightharpoonup u \text{ weakly in } L^2(0, T; V) \quad (78) \quad \boxed{\text{convgsd}}$$

Furthermore, note that $\tilde{u}_k(t - \tau_k) - \tilde{w}_k(t - \tau_k) \in V_t$ for every $t \in [0, T]$, and

$$\tilde{u}_k(\cdot - \tau_k) - \tilde{w}_k(\cdot - \tau_k) \rightharpoonup u - w \text{ weakly in } L^2(0, T; V). \quad (79)$$

Since the linear subspace $\{v \in L^2(0, T; V) \mid v(t) \in V_t \text{ for a.e. } t \in [0, T]\}$ is strongly closed, it is also weakly closed in $L^2(0, T; V)$. Therefore $u(t) \in V_t + w(t)$ for a.e. $t \in [0, T]$. Moreover for every $t \in (0, T]$ there exist an increasing sequence $t_i \in [0, T]$ converging to t such that $u(t_i) - w(t_i) \in V_{t_i}$ for every i . Thanks to (65) we know that $t \mapsto u(t) - w(t)$ is a strongly continuous map with values in V , and we obtain $u(t) - w(t) \in V_t$ for every $t \in (0, T]$. Together with the initial condition $u(0) = u^0 \in V_0$ we obtain (29). Moreover thanks to (75) and (76) we are in position to apply Lemma 2.12 to the function $u - w$ and hence to deduce that for a.e. $t \in [0, T]$

$$\frac{\dot{u}(t+h) - \dot{u}(t)}{h} \rightharpoonup \ddot{u}(t) \text{ weakly in } V_t^* \text{ as } h \rightarrow 0^+. \quad (80) \quad \boxed{\text{wconttd}}$$

Now we want to show that (33) holds for a.e. $t \in [0, T]$ for every $\phi \in V_t$. We claim that there exists a negligible set $W \subset [0, T]$ such that for $s \in D$ and for all $\phi \in V_s$, we have

$$\langle \ddot{u}(t), \phi \rangle_s + \langle \mathbb{C} \mathcal{E}u(t), \mathcal{E}\phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E}\dot{u}(t), \mathcal{E}\phi \rangle_{H_n} = \langle f(t), \phi \rangle_s, \quad (81) \quad \boxed{\text{claim}}$$

for every $t \in (s, T] \setminus W$.

To prove the claim, first we fix $s \in D$ and $\phi \in V_s$. Using (61) we have for a.e. $t > s$

$$\langle \dot{v}_k(t), \phi \rangle_s + \langle \mathbb{C} \mathcal{E} \tilde{u}_k(t), \mathcal{E} \phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E} \dot{u}_k(t), \mathcal{E} \phi \rangle_{H_n} = \langle f_k(t), \phi \rangle_s. \quad (82)$$

disped2d

Hence we have also

$$\int_s^T (\langle \dot{v}_k(t), \phi \rangle_s + \langle \mathbb{C} \mathcal{E} \tilde{u}_k(t), \mathcal{E} \phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E} \dot{u}_k(t), \mathcal{E} \phi \rangle_{H_n} - \langle f_k(t), \phi \rangle_s) dt = 0. \quad (83)$$

idispded

By construction $f_k \rightarrow f$ strongly in $L^2(0, T; V^*)$. We know that $\dot{v}_k \rightarrow \dot{v}$ weakly in $L^2(s, T; V_s^*)$ by (122). Since $\dot{u} = v$ in $W^{1,2}(s, T; V_s^*)$, we also have that $\ddot{u} = \dot{v}$ in $L^2(s, T; V_s^*)$. Using also (65) and (78), we can pass to the limit in (83) to have

$$\int_s^T (\langle \ddot{u}(t), \phi \rangle_s + \langle \mathbb{C} \mathcal{E} u(t), \mathcal{E} \phi \rangle_{H_n} + \langle \mathbb{B} \mathcal{E} \dot{u}(t), \mathcal{E} \phi \rangle_{H_n} - \langle f(t), \phi \rangle_s) dt = 0. \quad (84)$$

azero

By (82) we deduce that the integrand in (84) is zero for a.e. $t > s$. Since V_s is separable, the set N_s of $t > s$ for which (82) does not hold can be taken independent of ϕ . We set W to be the union over $s \in D$ of the sets N_s , so that W also has measure zero. It follows that for every $s \in D$ and for every $t \in (s, T] \setminus W$ (81) holds, and this shows the claim.

Using Lemma 2.12 it follows that for a.e. t and for every $\phi \in V_t$, there exist $s_i \nearrow t$ with $s_i \in D$ and $\phi_i \in V_{s_i}$, such that $\phi_i \rightarrow \phi$ strongly in V_t . Now note that if t belongs also to $(0, T] \setminus W$, by our previous claim we have

$$\begin{aligned} \langle \ddot{u}(t), \phi_i \rangle_t + \langle \mathbb{C} \mathcal{E} u(t), \mathcal{E} \phi_i \rangle_{H_n} + \langle \mathbb{B} \mathcal{E} \dot{u}(t), \mathcal{E} \phi_i \rangle_{H_n} - \langle f(t), \phi_i \rangle_t \\ = \langle \ddot{u}(t), \phi_i \rangle_{s_i} + \langle \mathbb{C} \mathcal{E} u(t), \mathcal{E} \phi_i \rangle_{H_n} + \langle \mathbb{B} \mathcal{E} \dot{u}(t), \mathcal{E} \phi_i \rangle_{H_n} - \langle f(t), \phi_i \rangle_{s_i} = 0. \end{aligned} \quad (85)$$

The convergence of the ϕ_i to ϕ gives (33).

Since by construction

$$f_k \rightarrow f \text{ strongly in } L^2(0, T; V^*), \quad (86)$$

$$w_k \rightarrow w \text{ strongly in } W^{1,2}(0, T; H^1(\Omega)^n), \quad (87)$$

$$\dot{w}_k \rightarrow \dot{w} \text{ strongly in } H \text{ for every } t \in [0, T], \quad (88)$$

$$\dot{z}_k \rightarrow L^2(0, T; H) \text{ strongly in } L^2(0, T; H), \quad (89)$$

using also (65) and (78), passing to the limit as $k \rightarrow \infty$ in (55), we obtain (39) by lower semicontinuity.

To prove (34), it is enough to show that for every sequence t_i of Lebesgue points for $t \mapsto \|\dot{u}(t)\|_H^2 + \|\mathcal{E}u(t)\|_{H_n^c}$ with $t_i \rightarrow 0$, then

$$\dot{u}(t_i) \rightarrow u^1 \text{ strongly in } H, \quad (90)$$

incondmrd

By (30) \dot{u} belongs to $W^{1,2}(0, T; V_0^*)$, so that, if $t_i \rightarrow 0$, then $\dot{u}(t_i) \rightarrow u^1$ in V_0^* . Since $\dot{u}(t_i)$ is bounded in H by (28) and $V_0 \subset H$ is dense, it follows that $\dot{u}(t_i) \rightarrow u^1$ weakly in H . Since $u \in W^{1,2}(0, T; H)$ then $u(t_i) \rightarrow u^0$ strongly in H , and since $\mathcal{E}u(t_i)$ is bounded in H_n , it follows that $\mathcal{E}u(t_i) \rightarrow \mathcal{E}u^0$ weakly in H_n . Therefore (90) is equivalent to

$$\begin{aligned} \limsup_{i \rightarrow \infty} \|\dot{u}(t_i)\|_H^2 + \|\mathcal{E}u^0\|_{H_n^c} &\leq \limsup_{i \rightarrow \infty} \|\dot{u}(t_i)\|_H^2 + \liminf_{i \rightarrow \infty} \|\mathcal{E}u(t_i)\|_{H_n^c} \\ &\leq \limsup_{i \rightarrow \infty} \|\dot{u}(t_i)\|_H + \|\mathcal{E}u(t_i)\|_{H_n^c} \\ &\leq \|u^1\|_H + \|\mathcal{E}u^0\|_{H_n^c}. \end{aligned} \quad (91)$$

lsimrd

By (28) and (39) there exists a constant C such that

$$\|\dot{u}(t)\|_H^2 + \|\mathcal{E}u(t)\|_{H_n^c}^2 \leq \|u^1\|_H^2 + \|\mathcal{E}u^0\|_{H_n^c}^2 + o(1) \quad (92)$$

enine1d

for a.e. $t \in [0, T]$. Inequality (92) holds for all Lebesgue points of $t \mapsto \|\dot{u}(t)\|_H^2 + \|\mathcal{E}u(t)\|_{H_n^c}^2$, in particular for $t = t_i$, so that the last inequality in (91) follows from (92). \square

contoz

Proposition 3.4. *Let u be the function given by Theorem 4.2, then u satisfies condition (32) and the energy balance (38) for every t Lebesgue point of $\|\dot{u}(\cdot)\|_H^2$.*

Proof. Let w be the Dirichlet boundary condition considered in Definition 3.1. We note that for every $h \in (0, T)$ and for a.e. $t \in [0, T]$ the functions $u(t) - u(t-h) - (w(t) - w(t-h)) \in V_t$. Hence if we define $z(t) := u(t) - w(t)$ we can test equation (33) with $\frac{z(t) - z(t-h)}{h}$ and integrate on (h, T) to get

$$\begin{aligned} & \int_h^t \left\langle \ddot{u}(\tau), \frac{z(\tau) - z(\tau-h)}{h} \right\rangle_\tau d\tau + \int_h^t \left\langle \mathbb{C} \mathcal{E}u(\tau), \frac{\mathcal{E}z(\tau) - \mathcal{E}z(\tau-h)}{h} \right\rangle_{H_n} d\tau \\ & + \int_h^t \left\langle \mathbb{B} \mathcal{E}\dot{u}(\tau), \frac{\mathcal{E}z(\tau) - \mathcal{E}z(\tau-h)}{h} \right\rangle_{H_n} d\tau \\ & = \int_h^t \left\langle f(\tau), \frac{z(\tau) - z(\tau-h)}{h} \right\rangle_\tau d\tau + \int_h^t \left\langle F(\tau), \frac{z(\tau) - z(\tau-h)}{h} \right\rangle_\Theta d\tau. \end{aligned} \quad (93)$$

Since $u \in W^{1,2}(0, T; V)$ and $w \in W^{2,2}(0, T; H) \cap W^{1,2}(0, T; V)$, we take the limit as $h \rightarrow 0^+$ on both side of the previous equality

$$\begin{aligned} & \lim_{h \rightarrow 0^+} \int_h^t \left\langle \ddot{u}(\tau), \frac{z(\tau) - z(\tau-h)}{h} \right\rangle_\tau d\tau + \int_0^t \langle \mathbb{C} \mathcal{E}\dot{u}(\tau), \mathcal{E}\dot{z}(\tau) \rangle_{H_n} d\tau \\ & + \int_h^t \langle \mathbb{C} \mathcal{E}u(\tau), \mathcal{E}\dot{z}(\tau) \rangle_{H_n} d\tau = \int_0^t \langle f(\tau), \dot{z}(\tau) \rangle_\tau d\tau + \int_0^t \langle F(\tau), \dot{z}(\tau) \rangle_\Theta d\tau \end{aligned} \quad (94)$$

enbalp

In order to compute the limit of the first term in the left hand-side of (94), we use Lemma 2.14 to write

$$\begin{aligned} & \int_h^t \left\langle \ddot{u}(\tau), \frac{z(\tau) - z(\tau-h)}{h} \right\rangle_\tau d\tau = \left\langle \dot{u}(t), \frac{z(t) - z(t-h)}{h} \right\rangle_H - \left\langle \dot{u}(h), \frac{z(h) - z(0)}{h} \right\rangle_H \\ & - \int_h^t \left\langle \dot{u}(\tau), \frac{\dot{z}(\tau) - \dot{z}(\tau-h)}{h} \right\rangle_H d\tau \end{aligned} \quad (95)$$

exptt

Since t is a Lebesgue point of $\|\dot{u}(\cdot)\|_H^2$ and u satisfies the initial condition (34) we have

$$\lim_{h \rightarrow 0^+} \left\langle \dot{u}(t), \frac{u(t) - u(t-h)}{h} \right\rangle_H = \|\dot{u}(t)\|_H^2 \quad \text{and} \quad \lim_{h \rightarrow 0^+} \left\langle \dot{u}(h), \frac{u(h) - u(0)}{h} \right\rangle_H = \|u^1\|_H^2.$$

Moreover using the identity $\langle \dot{u}(\tau), \dot{u}(\tau) - \dot{u}(\tau-h) \rangle_H = \frac{1}{2} \|\dot{u}(\tau-h)\|_H^2 - \frac{1}{2} \|\dot{u}(\tau)\|_H^2 + \frac{1}{2} \|\dot{u}(\tau) - \dot{u}(\tau-h)\|_H^2$, we can write

$$\begin{aligned} & \int_h^t \left\langle \dot{u}(\tau), \frac{\dot{u}(\tau) - \dot{u}(\tau-h)}{h} \right\rangle_H d\tau = \frac{1}{2h} \int_t^{t+h} \|\dot{u}(\tau)\|_H^2 d\tau - \frac{1}{2h} \int_0^h \|\dot{u}(\tau)\|_H^2 d\tau \\ & + \frac{1}{2h} \int_h^t \|\dot{u}(\tau) - \dot{u}(\tau-h)\|_H^2 d\tau. \end{aligned}$$

Again using condition (34) and the fact that t is a Lebesgue point of $\|\dot{u}(\cdot)\|_H^2$, we can write

$$\begin{aligned} \lim_{h \rightarrow 0^+} \int_h^t \left\langle \dot{u}(\tau), \frac{\dot{u}(\tau) - \dot{u}(\tau - h)}{h} \right\rangle_H d\tau &= \frac{1}{2} \|\dot{u}(t)\|_H^2 - \frac{1}{2} \|u^1\|_H^2 \\ &+ \lim_{h \rightarrow 0^+} \frac{1}{2} \int_h^t \frac{\|\dot{u}(\tau) - \dot{u}(\tau - h)\|_H^2}{h} d\tau, \end{aligned} \quad (96)$$

Finally using $w \in W^{2,2}(0, T; H)$ we can take the limit on both side of (95)

$$\begin{aligned} \lim_{h \rightarrow 0^+} \int_h^t \left\langle \ddot{u}(\tau), \frac{z(\tau) - z(\tau - h)}{h} \right\rangle_\tau d\tau &= \frac{1}{2} \|\dot{u}(t)\|_H^2 - \frac{1}{2} \|u^1\|_H^2 \\ &+ \lim_{h \rightarrow 0^+} \frac{1}{2} \int_h^t \frac{\|\dot{u}(\tau) - \dot{u}(\tau - h)\|_H^2}{h} d\tau + \frac{1}{2} \langle \dot{u}(t), \dot{w}(t) \rangle_H \\ &- \frac{1}{2} \langle u^1, \dot{w}(0) \rangle_H + \frac{1}{2} \int_0^t \langle \dot{u}(\tau), \ddot{w}(\tau) \rangle_H d\tau. \end{aligned} \quad (97) \quad \boxed{\text{lswtfih}}$$

Putting together (94) with (97) we obtain for every Lebesgue point t of $\|\dot{u}(\cdot)\|_H^2$

$$\begin{aligned} \frac{1}{2} \|\dot{u}(t)\|_H^2 + \frac{1}{2} \|\mathcal{E}u(t)\|_{H_n^c}^2 + \int_0^t \|\mathcal{E}\dot{u}(\tau)\|_{H_n^B}^2 d\tau - \lim_{h \rightarrow 0^+} \frac{1}{2} \int_h^t \frac{\|\dot{u}(\tau) - \dot{u}(\tau - h)\|_H^2}{h} d\tau \\ = \frac{1}{2} \|u^1\|_H^2 + \frac{1}{2} \|\mathcal{E}u^0\|_{H_n^c}^2 + \mathcal{W}_{load}(u; 0, t) + \mathcal{W}_{bdry}(u; 0, t). \end{aligned} \quad (98) \quad \boxed{\text{citrem}}$$

But since we already know u satisfies the energy inequality (39), we immediately conclude that both condition (32) and the energy balance (38) hold. \square

remfint

Remark 3.5. At each time t we expect that any energy increment for the solution $u(t)$, is due to the external forces. By identity (98), which is true for every u satisfying (28)-(31) and (33), the requirement (32) i.e.

$$\lim_{h \rightarrow 0^+} \frac{1}{2} \int_h^t \frac{\|\dot{u}(\tau) - \dot{u}(\tau - h)\|_H^2}{h} d\tau = 0$$

is natural in term of energy balance, and allow us to deal with solutions whose energy behavior is consistent from a physical point of view.

Remark 3.6. Since $\dot{u}(t) - \dot{w}(t) \in V_t$ for a.e. $t \in [0, T]$, we can use it as test function in (33) and integrate on $(0, t)$ to obtain

$$\begin{aligned} \int_0^t \langle \ddot{u}(\tau), \dot{u}(\tau) \rangle_\tau d\tau + \frac{1}{2} \|\mathcal{E}u(t)\|_{H_n^c}^2 + \int_0^t \|\mathcal{E}\dot{u}(\tau)\|_{H_n^B}^2 d\tau \\ = \frac{1}{2} \|\mathcal{E}u^0\|_{H_n^c}^2 + \mathcal{W}_{load}(u; 0, t) + \mathcal{W}_{bdry}(u; 0, t). \end{aligned}$$

Comparing this last identity with the energy balance (38), we have for a.e. $t \in (0, T)$ that

$$\frac{1}{2} \|\dot{u}(t)\|_H^2 - \frac{1}{2} \|u^1\|_H^2 = \int_0^t \langle \ddot{u}(\tau), \dot{u}(\tau) \rangle_\tau d\tau,$$

and since $\tau \mapsto \langle \ddot{u}(\tau), \dot{u}(\tau) \rangle_\tau \in L^1(0, T)$ we deduce that $\|\dot{u}(\cdot)\|_H^2 \in W^{1,1}(0, T)$.

Putting together Theorem 3.2 and Proposition 3.4, we deduce the existence of a solution u to the damped system of elastodynamic. Moreover using (98) we can also obtain the uniqueness of weak solutions considered in Definition 3.1. This is the content of the next theorem.

exun

Theorem 3.7 (Existence and uniqueness). *Under hypothesis of Theorem 3.2 there exists a unique solution u considered in Definition 3.1. Moreover u satisfies the energy balance*

$$\begin{aligned} & \frac{1}{2} \|\dot{u}(t)\|_H^2 + \frac{1}{2} \|\mathcal{E}u(t)\|_{H_n^c}^2 + \frac{1}{2} \int_0^t \frac{1}{2} \|\mathcal{E}\dot{u}(\tau)\|_{H_n^{\mathbb{E}}}^2 d\tau = \\ & = \frac{1}{2} \|u^1\|_H^2 + \frac{1}{2} \|\mathcal{E}u^0\|_{H_n^c}^2 + \mathcal{W}_{load}(u; 0, t) + \mathcal{W}_{bdry}(u; 0, t) \end{aligned} \quad (99)$$

enbald1

for a.e. $t \in [0, T]$.

Proof. The existence of a solution satisfying (99) is simply a consequence of Theorem 3.2 and Proposition 3.4.

To show uniqueness, we notice that (28)-(33) are all preserved under linear combinations. Therefore, the difference v between two solutions is a solution with Dirichlet and Neumann homogeneous conditions, with forcing term $f = 0$, and satisfying $v(0) = 0$ and

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_0^h \|\dot{v}(t)\|_h^2 dt = 0.$$

Moreover using the same argument as in Proposition, since (32) holds for v , we have

$$\frac{1}{2} \|\dot{v}(t)\|_H^2 + \frac{1}{2} \|\mathcal{E}v(t)\|_{H_n^c}^2 + \frac{1}{2} \int_0^t \frac{1}{2} \|\mathcal{E}\dot{v}(\tau)\|_{H_n^{\mathbb{E}}}^2 d\tau = 0,$$

for a.e. $t \in [0, T]$. Therefore $\dot{v}(t) = 0$ a.e. on $[0, T]$. Since $v \in W^{1,\infty}(0, T; H)$ and $v(0) = 0$, we conclude $v(t) = 0$ a.e. on $[0, T]$. \square

Finally, one can also prove that the energy balance (38) holds for every $t \in [0, T]$ and that the map $t \mapsto \dot{u}(t)$ is strongly continuous in H . For the proof of this result we refer to [8, Lemma 3.10].

notproved

Proposition 3.8. *Under the assumptions of Theorem 3.2, let u be the weak solution of the damped wave equation considered in Definition (3.1), with initial conditions $u(0) = u^0$ and (34). Then $t \mapsto \dot{u}(t)$ is continuous from $[0, T]$ to H and the energy balance (38) holds for every $t \in [0, T]$.*

4. THE UNDAMPED SYSTEM OF ELASTODYNAMICS

In this section we study weak solutions of the undamped system of elastodynamics

$$\ddot{u}(t) - \operatorname{div}[\mathbb{C}\mathcal{E}u(t)] = f(t), \quad (100)$$

sysedu

on a cracking domain.

ds

Definition 4.1. Assume (a), (b), (c) and (d). With the notation introduced in Section 2, let $f \in W^{1,2}(0, T; V^*)$, let $w \in W^{2,2}(0, T; H) \cap W^{1,2}(0, T; H^1(\Omega)^n)$ and let $F \in W^{1,2}(0, T; N_{\Theta})$ where Θ is the function relative to the crack set Γ given by Theorem 2.3. We say that u is a weak solution of (1) on the time dependent domain $t \mapsto \Omega \setminus \Gamma(t)$ with Dirichlet boundary condition $w(t)$ on $\partial_D \Omega$, Neumann boundary condition $F(t)$ on $\partial_N \Omega$, and homogeneous Neumann boundary conditions on $\Gamma(t)$,

if

$$u \in L^\infty(0, T; V) \cap W^{1, \infty}(0, T; H). \quad (101) \quad \boxed{\text{ds2}}$$

$$\text{For every } t \in [0, T] \ u(t) - w(t) \in V_t. \quad (102) \quad \boxed{\text{ds3}}$$

$$\text{For every } s \in [0, T] \ u \in W^{2,2}(s, T; V_s^*) \text{ and} \quad (103) \quad \boxed{\text{ds4}}$$

$$\|P_{st}\ddot{u}(t)\|_s^* \leq g(t) \text{ for a.e. } t \in (s, T), \text{ for some } g \in L^2(0, T). \quad (104) \quad \boxed{\text{ds5}}$$

For a.e. $t \in [0, T]$

$$\langle \ddot{u}(t), \phi \rangle_t + \langle \mathcal{C}\mathcal{E}u(t), \mathcal{E}\phi \rangle_{H_n} - \langle F(t), \text{Tr}(\phi) \rangle_\Theta = \langle f(t), \phi \rangle_t, \text{ for every } \phi \in V_t \quad (105) \quad \boxed{\text{ds7}}$$

where $\ddot{u}(t)$ is the one given by Definition 2.13.

Given $u^0 \in V_0$ and $u^1 \in H$, since $t \mapsto u(t)$ is strongly continuous in H the initial value for u^0 is well defined as element of H . Moreover we are able to prescribe the initial conditions respectively for $\mathcal{E}u(0)$ and $\dot{u}(0)$ asking

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_0^h \|\mathcal{E}u(t) - \mathcal{E}u^0\|_{H_n}^2 dt = 0, \quad (106) \quad \boxed{\text{incond1}}$$

and

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_0^h \|\dot{u}(t) - u^1\|_H^2 dt = 0. \quad (107) \quad \boxed{\text{incond}}$$

Since in this case, $\dot{u}(t)$ is in general only an element in H , we need to consider also a weakened formulation of the work due to the Neumann boundary conditions. More precisely we define

$$\begin{aligned} \mathcal{W}_{bdry}^N(u; t_1, t_2) := & \int_{\partial_N \Omega} F(t_2) \cdot u(t_2) d\mathcal{H}^{n-1} - \int_{\partial_N \Omega} F(t_1) \cdot u(t_1) d\mathcal{H}^{n-1} \\ & - \int_{t_1}^{t_2} \left(\int_{\partial_N \Omega} \dot{F}(t) \cdot u(t) d\mathcal{H}^{n-1} \right) dt \end{aligned} \quad (108)$$

for every time interval $[t_1, t_2] \subset [0, T]$.

This is the main result:

mres

Theorem 4.2. *Assume (a), (b), (c) and (d). Let f , w and F be as in Definition 4.1. Then, given two initial conditions $u^0 \in V_0$ and $u^1 \in H$, there exists a solution u of (1) with initial conditions (106) and (107). Moreover u satisfies the energy inequality*

$$\frac{1}{2} \|\dot{u}(t)\|_H^2 + \frac{1}{2} \|\mathcal{E}u(t)\|_{H_n^c}^2 \leq \frac{1}{2} \|u^1\|_H^2 + \frac{1}{2} \|\mathcal{E}u^0\|_{H_n^c}^2 + \mathcal{W}_{load}(u; 0, t) + \mathcal{W}_{bdry}(u; 0, t) \quad (109) \quad \boxed{\text{enine}}$$

for a.e. $t \in [0, T]$.

Proof. Since the argument is similar to the one given for Theorem 4.2, we simply give a sketch of the proof.

For $k \in \mathbb{N}$, we set $\tau_k := T/k$ and $t_k^j := j\tau_k$. For $j = 1, 2, \dots, k$ we define $f_k^j \in V^*$ and $w_k^j \in H^1(\Omega)^n$ by

$$f_k^j := f(t_k^j), \quad w_k^j := w(t_k^j), \quad (110)$$

using that $f \in W^{1,2}(0, T; V^*)$ and $w \in W^{1,2}(0, T; H^1(\Omega)^n)$, so f and w are well defined elements of V^* and $H^1(\Omega)^n$, respectively, for every $t \in [0, T]$. Inductively we define u_k^j for $j = -1, 0, \dots, k$ by the following:

$$u_k^0 := u^0, \quad u_k^{-1} := u^0 - \tau_k u^1; \quad (111) \quad \boxed{\text{condin}}$$

then, for $j = 0, 1, \dots, k-1$, the function u_k^{j+1} is the minimizer in $V_{t_k^{j+1}} + w_k^j$ of

$$u \mapsto \left\| \frac{u - u_k^j}{\tau_k} - \frac{u_k^j - u_k^{j-1}}{\tau_k} \right\|_H^2 + \|\mathcal{E}u\|_{H_n^c}^2 - 2\langle f_k^j, u \rangle_{t_k^{j+1}}. \quad (112)$$

Then if we define $u_k, \tilde{u}_k, w_k, \tilde{w}_k, v_k$, and z_k as in (49)-(53), then proceeding exactly as in 3.2, we deduce the following bounds

$$\mathcal{E}u_k(t) \text{ and } \mathcal{E}\tilde{u}_k(t) \text{ are bounded in } H_n \text{ uniformly in } t \text{ and } k, \quad (113)$$

bpd1

$$\dot{u}_k(t) \text{ and } v_k(t) \text{ are bounded in } H \text{ uniformly in } t \text{ and } k \quad (114)$$

bpd3

$$u_k(t) \text{ is bounded in } V \text{ uniformly in } t \text{ and } k. \quad (115)$$

bpd4

Furthermore, using the Euler equation for u_k^{j+1} we can write for all $t \in (t_k^j, t_k^{j+1})$

$$\langle \dot{v}_k(t), \phi \rangle_H + \langle \mathbb{C} \mathcal{E}\tilde{u}_k(t), \mathcal{E}\phi \rangle_{H_n} = \langle f_k(t), \phi \rangle_{t_k^{j+1}} \quad (116)$$

dispde1

for every $\phi \in V_{t_k^{j+1}}$. The last equation leads us to write for all $t \in (t_k^j, t_k^{j+1})$

$$\|\dot{v}_k(t)\|_{t_k^{j+1}}^* \leq \|\mathcal{E}\tilde{u}_k(t)\|_{H_n^c} + \|f_k(t)\|_{t_k^{j+1}}^*. \quad (117)$$

bpd5

In particular fix $s \in [0, T]$, then for every $t_1, t_2 \in [s, T]$ with $t_1 < t_2$, we have

$$\int_{t_1}^{t_2} \|P_{st}\dot{v}_k(t)\|_s^* dt \leq \int_{t_1}^{t_2} (\|\mathcal{E}\tilde{u}_k(t)\|_{H_n^c} + \|f_k(t)\|_s^*) dt. \quad (118)$$

acest

Again following exactly 3.2 we deduce that up to subsequences

$$u_k \rightharpoonup u, \text{ weakly in } W^{1,2}(0, T; H), \quad (119)$$

estlem3

$$\tilde{u}_k \rightharpoonup u, \text{ weakly in } L^2(0, T; V) \quad (120)$$

estlem6

$$v_k \rightharpoonup v \text{ weakly in } L^2(0, T; H), \quad (121)$$

estlem2

$$v_k \rightharpoonup v \text{ weakly in } W^{1,2}(s, T; V_s^*). \quad (122)$$

estlem1

Moreover $u \in L^\infty(0, T; V) \cap W^{1,\infty}(0, T; H)$, $\dot{u}(t) = v(t)$ a.e. on $[0, T]$, and

$$\|P_{st}\ddot{u}(t)\|_t^* \leq M + \|f(t)\|_s^*, \quad (123)$$

for every $s \in [0, T]$ and for a.e. $t > s$.

Now the proof that u is a solution, and that satisfies the energy inequality (109) proceeds as in the damped case. Finally, it remains to prove that u satisfies the initial conditions (106) (107). It is enough to show that for every sequence t_i of Lebesgue points for $t \mapsto \|\dot{u}(t)\|_H^2 + \|\mathcal{E}u(t)\|_{H_n^c}^2$ and $t_i \rightarrow 0$, then

$$\dot{u}(t_i) \rightarrow u^1 \text{ strongly in } H, \quad (124)$$

incondmr

and

$$\mathcal{E}u(t_i) \rightarrow \mathcal{E}u^0 \text{ strongly in } H_n. \quad (125)$$

Again this can be achieved following the same argument as in the damped case \square

REFERENCES

- ada** [1] Adams R.A.: *Sobolev Spaces* Academic Press, 1975.
- afp** [2] Ambrosio L., Fusco N., Pallara D.: *Functions of bounded variation and free discontinuity problems*. Oxford University Press, Oxford, 2000.
- cap** [3] Caponi M., Lucardesi I., Tasso E.: *Energy dissipation balance of a smooth moving crack*. Preprint SISSA 31/2018/MATE.
- iur1** [4] Chambolle A., Conti S., Iurlano I.: *Approximation of functions with small jump sets and existence of strong minimizers of Griffith's energy*. Preprint available on: <http://cvgmt.sns.it/paper/3599/>.
- iur2** [5] Conti S., Focardi M., Iurlano F.: *Existence of strong minimizers for the Griffith static fracture model in dimension two*. Annales de l'Institut Henri Poincaré C, Analyse non linéaire. Elsevier Masson, 2018.
- dalmaso1** [6] Dal Maso G., *Generalised functions of bounded deformation*, J. Eur. Math. Soc. (JEMS), to appear. Preprint available on: <http://cvgmt.sns.it/paper/1704/>.
- dalmaso4** [7] Dal Maso G., Francfort G.A., Toader R.: *Quasistatic Crack Growth in Finite Elasticity*. Preprint SISSA, Trieste, 2004, <http://www.sissa.it/fa/>.
- dalmaso2** [8] Dal Maso G., Larsen C. J.: *Existence for wave equations in domains with arbitrary growing cracks*. Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 22, 2011, 387408.
- dalmaso7** [9] Dal Maso G., Larsen C. J., Toader R.: *Existence for constrained dynamic Griffith fracture with a weak maximal dissipation condition*. Journal of the Mechanics and Physics of Solids 95, 2016, 697-707.
- dalmaso5** [10] Dal Maso G., Lazzaroni G.: *Quasistatic crack growth in finite elasticity with non-interpenetration*. Annales de l'Institut Henri Poincaré (C) Non Linear Analysis. Vol. 27. No. 1. Elsevier Masson, 2010.
- dalmaso6** [11] Dal Maso G., Toader R.: *A Model for the Quasi-Static Growth of Brittle Fractures: Existence and Approximation Results*. Archive for Rational Mechanics and Analysis 162.2, 2002, 101-135.
- eva** [12] Evans L.,Gariepy R.F.: *Measure theory and fine properties of functions*. CRC Press, Boca Raton, 1992.
- fed** [13] Federer H.: *Geometric measure theory*. Springer-Verlag, Berlin, 1969.
- fra1** [14] Francfort, G.A., Larsen C.J.: *Existence and convergence for quasi-static evolution in brittle fracture*. Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences 56.10, 2003, 1465-1500.
- fre** [15] Freund L. B.: *Dynamic Fracture Mechanics*. Cambridge Univ. Press, New York, 1990.
- lar** [16] Larsen C. J.: *Models for dynamic fracture based on Griffith's criterion*. In IUTAM Symposium on Variational Concepts with Applications to the Mechanics of Materials (Klaus Hackl, ed.) Springer, 2010, 131-140.
- ol** [17] Oleinik O.A., Shamaev A.S., Yosifan G.A.: *Mathematical Problems in Elasticity and Homogenization*. Studies in Mathematics and Applications, North Holland, 1992.
- tax** [18] Tasso E.: *On the continuity of the trace operator in $GSBV(\Omega)$ and $GSBD(\Omega)$* . Preprint SISSA 39/2018/MATE.
- tem** [19] Temam R., *Mathematical problems in plasticity*. Gauthier-Villars, Paris, 1985. Translation of *Problèmes mathématiques en plasticité*. Gauthier-Villars, Paris, 1983.

(Emanuele Tasso) SISSA, VIA BONOMEA 265, 34136 TRIESTE, ITALY
 E-mail address: etasso@sissa.it